

# Class 24 Section 5.3 I ~~II~~

Note: The monotonicity of  $f(x)$  and its concavity are not related. (why?)

ex. Find the interval(s) where  $f(x) = \frac{x-6}{x+2}$  is increasing or decreasing and also its concavity.

Solution: We need both  $f'(x)$  and  $f''(x)$  here.

•  $f'(x) = \frac{1(x+2) - 1(x-6)}{(x+2)^2} = \frac{8}{(x+2)^2}$ . The domain is  $D = \{x \in \mathbb{R} \mid x \neq -2\}$  and  $f'(x) > 0$  on all of  $D$ .  $f(x)$  is increasing on all of  $D$ .


•  $f''(x) = \frac{d}{dx} \left[ \frac{8}{(x+2)^2} \right] = \frac{0(x+2)^2 - 8 \cdot 2(x+2)}{(x+2)^4} = \frac{-16(x+2)}{(x+2)^4}$

$= \frac{-16}{(x+2)^3}$ . Here  $f''(x) > 0$  when  $x < -2$

hence concave up on  $(-\infty, -2)$ ,

and concave down on  $(-2, \infty)$

since  $f''(x) < 0$  here.



So let  $f(x)$  be twice differentiable on some interval  $I = (a, b)$ , and at a pt  $c \in (a, b)$   $f'(c) = 0$  and  $f''(c) < 0$ . What is happening here? And when  $f'(c) = 0$  and  $f''(c) > 0$ ?


### Second Derivative Test for a local extremum

Suppose  $f$  is twice diff on an open interval, containing a pt  $c$ :


If  $f'(c) = 0$  and  $f''(c) < 0$ , then  $f(c)$  is a local max

If  $f'(c) = 0$  and  $f''(c) > 0$ , then  $f(c)$  is a local min.


And if both  $f'(c) = 0$  and  $f''(c) = 0$ ? Nothing.



$$f'(0) = f''(0) = 0$$



$$g'(0) = g''(0) = 0$$



$$h'(0) = h''(0) = 0$$

ex. Find the extreme of  $f(x) = xe^{-x}$  on the interval  $[0, 10]$ .

Solution: ~~Let~~ Since  $f'(x) = e^{-x} - xe^{-x} = (1-x)e^{-x}$  exists on all of  $(0, 10)$ , the extreme will occur either at an endpoint or at a place where  $f'(x) = 0 = (1-x)e^{-x}$ . Thus, the extreme will occur at  $x=0$ ,  $x=10$ , or  $x=1$ .

We could simply check the values here.

$$f(0) = 0, \quad f(10) = 10e^{-10}, \quad f(1) = \frac{1}{e}.$$

Hence minimum at  $x=0$ , and max at  $x=1$ .

We could also test the max at  $x=1$  by the

$$SDT: \quad f''(x) = -e^{-x} - (1-x)e^{-x} = (x-2)e^{-x}$$

Since  $f'(1) = 0$  and  $f''(1) = (1-2)e^{-1} < 0$  at  $x=1$ ,

we know by 2<sup>nd</sup> Der test that there is a local max at  $x=1$ .

