

110.302 Lecture 23: ~~XXXXXXXXXXXX~~ I

For $\vec{x}' = A_{2 \times 2} \vec{x}$,

(I) One can construct a slope field in \mathbb{R}^2 via matrix multiplication: For $\vec{x} \in \mathbb{R}^2$,

~~the tangent to the solution curve~~
 passing through $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is $A\vec{x}$

(a) $\vec{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\vec{x}' = \begin{bmatrix} 1 & 1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$.

(b) $\vec{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\vec{x}' = \begin{bmatrix} 1 & 1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \end{bmatrix}$

(c) $\vec{x} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $\vec{x}' = \begin{bmatrix} 1 & 1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

JUDE = 2D calculator (or similar) is helpful here.

(II) Solution curves are integral curves of the slope field:

(a) Given $c_1, c_2 \in \mathbb{R}$, the curve

$$\vec{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$$


is one of these curves.

③ Straight-line motion only occurs when one of c_1, c_2 is 0:

Let $c_1 = -2, c_2 = 0$.

$$\Rightarrow \vec{x}(t) = -2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t}$$

$$= \begin{bmatrix} -2 \\ -4 \end{bmatrix} e^{3t}$$



④ A copy of \mathbb{R}^2 with enough representative curves on it to give a good sense of solutions is called a phase portrait:

⑤ Solutions are called trajectories or orbits.

⑥ General long term behavior of trajectories can be read off easily from a phase portrait:

$$\text{Given } \vec{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} e^{-2t}$$


(II) (c) cont'd.

- If $c_1 = 0, c_2 \neq 0$, then $\vec{x}(t) = c_2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$

and $\lim_{t \rightarrow \infty} \vec{x}(t) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

How about $\lim_{t \rightarrow -\infty} \vec{x}(t)$?

We say it is unbounded.



Note: Solutions never touch nor cross here.

So since $\vec{x}(t) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ is an ~~an~~ equilibrium soln, no other solution actually reaches $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

- If $c_1 \neq 0, c_2 = 0$?

$\lim_{t \rightarrow \infty} \vec{x}(t)$ DNE or traj is unbounded


$\lim_{t \rightarrow -\infty} \vec{x}(t) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$.

- If $c_1 \neq 0$, and $c_2 \neq 0$? These are the curved lines in the portrait. For these, what can we say about the forward trajectory ($\lim_{t \rightarrow \infty} \vec{x}(t)$), or the backward one ($\lim_{t \rightarrow -\infty} \vec{x}(t)$)?

One answer? Given $\vec{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$

and $c_1 \neq 0, c_2 \neq 0$, then as t gets large ($t \rightarrow \infty$), $\vec{x}(t)$ looks more and more like

$c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t}$, and less and less like $c_2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$



How about in backward time?

III) Back to solution building via the properties

of A : For ~~\vec{x}~~ $\vec{x}' = \begin{bmatrix} 1 & 1 \\ 6 & 0 \end{bmatrix} \vec{x}$,

$$\vec{x}(t) = c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} e^{3t} + c_2 \begin{bmatrix} 1 \\ -3 \end{bmatrix} e^{-2t}$$

the eigenvalues of A (~~$\lambda_1 = 3, \lambda_2 = -2$~~), and representative eigenvectors ($\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$) are explicitly part of the solution.

Why?

Here, the eigenvalues and eigenvectors of a matrix A satisfy $A\vec{v} = \lambda\vec{v}$, or $(A - \lambda I)\vec{v} = \vec{0}$.

For this to have nontrivial solutions for \vec{v} ,

-
- λ must be an eigenvalue, and
 - $\det(A - \lambda I) = 0$.

$$\text{for } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \quad \det(A - \lambda I) = \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} = 0$$
$$= \lambda^2 - (a+d)\lambda + (ad - bc) = 0$$

This is called the characteristic eqn of A , and solutions can be

- ① real, distinct
- ② real and repeated
- ③ complex conjugates.

How to relate to solutions of $\dot{\vec{x}} = A\vec{x}$?

- Recall $\dot{x} = ax$ is solved by $x(t) = ce^{at}$
- Assume $\dot{\vec{x}} = A\vec{x}$ is also solved by exponentials. For $n=2$, $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$.

Assume $x_1(t) = c_1 e^{rt}$, $x_2(t) = c_2 e^{rt}$

$$\Rightarrow \vec{x}(t) = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} e^{rt}, \text{ and } \vec{x}' = r \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} e^{rt}$$

and then $\vec{x}' = A\vec{x}$ is $r \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} e^{rt} = A \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} e^{rt}$

or $r \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = A \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$, i.e. $r\vec{v} = A\vec{v}$.

And what do solutions to this look like??

r is an eigenvalue

\vec{v} is an eigenvector of A .

Notes ① This works equally well for $n > 2$.

② In the case where $A_{n \times n}$ is real and symmetric (i.e. when $a_{ij} = a_{ji}$)

\Rightarrow all eigenvalues are real and even if repeated, there is a full set of eigenvectors.

We have the following:

For $\dot{\vec{x}} = A_{n \times n} \vec{x}$, when all eigenvalues of A are real and distinct, then

$$\vec{x}(t) = c_1 \vec{v}_1 e^{r_1 t} + \dots + c_n \vec{v}_n e^{r_n t}$$

is the general soln, where \vec{v}_i is the c_i eigenvector of r_i , for A .