

# Chapter 2: Method of Separation of Variables

Fei Lu

Department of Mathematics, Johns Hopkins

Solution to the IBVP?

$$\partial_t u = \kappa \partial_{xx} u + Q(x, t), \quad \text{with } x \in (0, L), t > 0$$

$$u(x, 0) = f(x)$$

$$\text{BC: } u(0, t) = \phi(t), u(L, t) = \psi(t)$$

Section 2.5 Laplace's equation: solution examples

Section 2.5 Laplace's equation: qualitative properties

# Outline

Section 2.5 Laplace's equation: solution examples

Section 2.5 Laplace's equation: qualitative properties


# 1. Laplace's equation inside a rectangular

Consider the Laplace's equation

$$\nabla^2 u = \partial_{xx}u + \partial_{yy}u = 0, \quad 0 \leq x \leq L, 0 \leq y \leq H$$

$$u|_{\Gamma_1} = g_1(y); \quad u|_{\Gamma_2} = g_2(y);$$

$$u|_{\Gamma_3} = f_1(x); \quad u|_{\Gamma_4} = f_2(x);$$



- ▶ Equilibrium of the HE
- ▶ How to solve it? 1D:  $\partial_{xx}u = 0 \Rightarrow u(x) = c_1x + c_2$ .  
Separation of variables?  
Linear and homogeneous: PDE, BC

$$\nabla^2 u_1 = 0,$$

$$u_1|_{\Gamma_1} = g_1;$$

$$u_1|_{\Gamma_2} = 0;$$

$$u_1|_{\Gamma_3} = 0;$$

$$u_1|_{\Gamma_4} = 0;$$

$$\nabla^2 u_2 = 0,$$

$$u_2|_{\Gamma_1} = 0;$$

$$u_2|_{\Gamma_2} = g_2;$$

$$u_2|_{\Gamma_3} = 0;$$

$$u_2|_{\Gamma_4} = 0;$$

$$\nabla^2 u_3 = 0,$$

$$u_3|_{\Gamma_1} = 0;$$

$$u_3|_{\Gamma_2} = 0;$$

$$u_3|_{\Gamma_3} = f_1;$$

$$u_3|_{\Gamma_4} = 0;$$

$$\nabla^2 u_4 = 0,$$

$$u_4|_{\Gamma_1} = 0;$$

$$u_4|_{\Gamma_2} = 0;$$

$$u_4|_{\Gamma_3} = 0;$$

$$u_4|_{\Gamma_4} = f_2;$$

## Solve $u_1$ by Separation of Variables:

1. Seek solution  $u_1(x, y) = h(x)\phi(y)$ :

$$\nabla^2 u_1 = 0,$$

$$\frac{h''(x)}{h} = -\frac{\phi''(y)}{\phi} = \lambda$$

$$u_1|_{\Gamma_1} = g_1;$$

$$u_1|_{\Gamma_2} = 0;$$

$$u_1|_{\Gamma_3} = 0;$$

$$u_1|_{\Gamma_4} = 0;$$

2. Eigenvalue problem:

$$\phi''(y) = -\lambda\phi(y), \quad \phi(0) = \phi(H) = 0$$

$$\phi_n(y) = \sin\left(\frac{n\pi}{H}y\right), \quad \lambda_n = \left(\frac{n\pi}{H}\right)^2, \quad n = 1, 2, \dots,$$

3. Solve  $h$ :

$$h''(x) = \lambda h(x), \quad h(L) = 0$$

▶  $\lambda > 0$ :  $h(x) = c_1 e^{\sqrt{\lambda}x} + c_2 e^{-\sqrt{\lambda}x}$


$$h_n(x) = a_n \sinh(\sqrt{\lambda_n}(x - L))$$

4. Determine  $a_n$

$$u_1(x, y) = \sum_{n=1}^{\infty} a_n \sinh(\sqrt{\lambda_n}(x - L))\phi_n(y).$$

## 2.5.2 Laplace equation on a disk

$$\begin{aligned}\nabla^2 u &= 0, & (x, y) \in \text{Disk} \\ u|_{\Gamma} &= f\end{aligned}$$



$$x = r \cos \theta; \quad y = r \sin \theta$$

$$\nabla^2 u = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}, \quad 0 < r < a, -\pi < \theta < \pi$$

- ▶ BC:  $u(a, \pi) = u(a, -\pi)$ ;  $\partial_{\theta} u(a, \pi) = \partial_{\theta} u(a, -\pi)$   
 $u(a, \theta) = f(\theta)$ ;  $u(0, \theta) = ?$
- ▶ Separation of variables?  
linear homo: PDE, BC

$$\nabla^2 u = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0, \quad 0 < r < a, -\pi < \theta < \pi$$

BC:  $u(a, \pi) = u(a, -\pi)$ ;  $\partial_{\theta} u(a, \pi) = \partial_{\theta} u(a, -\pi)$   
 $u(a, \theta) = f(\theta)$ ;  $u(0, \theta) = ?$

- ▶ 1. Seek solution  $u(r, \theta) = G(r)\phi(\theta)$ :

$$\frac{r(rG')'}{G}(r) = -\frac{\phi''(\theta)}{\phi} = \lambda$$

- ▶ 2. EigenvalueP:

$$\phi''(\theta) = -\lambda\phi(\theta), \quad \phi(-\pi) = \phi(\pi); \phi'(-\pi) = \phi'(\pi)$$

- ▶ 3.  $G(r)$ :  $\frac{r(rG')'}{G} = \lambda_n$ ; BC?  $G(0)$  is bounded

- ▶ 4. Solution:  $\lambda_n = n^2$ ,  $\phi_n = \cos(n\theta), \sin(n\theta)$ ,  $n = 0, 1, \dots$

$$r^2 G'' + rG' - n^2 G = 0 \Rightarrow \text{(Euler's method:)} G(r) = r^p \text{ or } \ln r$$

$$u(r, \theta) = A_0 + \sum_{n=1}^{\infty} [A_n r^n \cos(n\theta) + B_n r^n \sin(n\theta)]$$

**\*2.5.3.** Solve Laplace's equation *outside* a circular disk ( $r \geq a$ ) subject to the boundary condition

(a)  $u(a, \theta) = \ln 2 + 4 \cos 3\theta$

(b)  $u(a, \theta) = f(\theta)$

You may assume that  $u(r, \theta)$  remains finite as  $r \rightarrow \infty$ .

$$\nabla^2 u = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial u}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0, \quad r > a, \quad -\pi < \theta < \pi$$

BC:  $u(a, \pi) = u(a, -\pi)$ ;  $\partial_\theta u(a, \pi) = \partial_\theta u(a, -\pi)$   
 $u(a, \theta) = g(\theta)$ ;  $\lim_{r \rightarrow \infty} u(r, \theta) < \infty$

# Outline

Section 2.5 Laplace's equation: solution examples

Section 2.5 Laplace's equation: qualitative properties



## 2.5.3 Qualitative properties

**Mean value property**  $u(P)$  is the average of  $u$  in  $\partial B_r(P) \subset D$

- ▶ The temperature at any point is the average of the temperature along any circle (inside domain) centered at the point.
- ▶ Example on disk:

$$u(0, \theta) = a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta) d\theta$$

**Maximum principle** In non-constant steady state the temperature cannot attain its maximum in the interior:

$$u(P) = \max_D u \Rightarrow P \in \partial D$$

## Wellposedness and uniqueness

Definition: a DE problem is well-posed if there **exists a unique** solution that *depends continuously* on the nonhomogeneous data.

### Theorem

$\nabla^2 u = 0$  on a smooth domain  $D$  with  $u|_{\partial D} = f(x)$  is well-posed.

### "Proof".

- ▶ Existence: physical intuition, for compatible  $f$ .  
solution on  $\mathbb{R}^d$ ; then constraint on  $D$  (Reading: Craig Evans, Partial Differential Equations)
- ▶ Continuous dependence on BC
  
- ▶ Uniqueness



**Solvability condition** For  $\nabla^2 u = 0$ , we have (Divergence theorem)

$$\oint \nabla u \cdot \mathbf{n} dS = \int \nabla^2 u dV = 0$$

- ▶ If Neumann BC  $-K_0 \nabla u \cdot \mathbf{n}$ , then we must have  $\oint \nabla u \cdot \mathbf{n} dS = 0$
- ▶ The net heat flow through the boundary must be zero for a steady state (with no source).

## Summary of Chp 2: Separation of variables

- ▶ Heat equation + BC + IC; Laplace +BC
- ▶ Linear + homogeneous  $\Rightarrow$  Principle of superposition
- ▶ Separation of variables

**Solution of HE + BC+ IC:**  $\partial_t u = \partial_{xx} u$ ,  $u(x, 0) = f(x)$

$$\begin{array}{ll} \text{Dirichlet } x \in (0, L) & f(x) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi}{L}x\right) \\ u(0, t) = u(L, t) = 0 & u(x, t) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi}{L}x\right) e^{-\lambda_n \kappa t} \end{array}$$

$$\begin{array}{ll} \text{Neuman } x \in (0, L) & f(x) = \sum_{n=0}^{\infty} A_n \cos\left(\frac{n\pi}{L}x\right) \\ \partial_x u(0, t) = \partial_x u(L, t) = 0 & u(x, t) = A_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi}{L}x\right) e^{-\lambda_n \kappa t} \end{array}$$

$$\begin{array}{ll} \text{Mixed } x \in (-L, L) & f(x) = A_0 + \sum_{n=1}^{\infty} \left( A_n \cos \frac{n\pi}{L}x + B_n \sin \frac{n\pi}{L}x \right) \\ \partial_x u(0, t) = \partial_x u(L, t) & u(x, t) = A_0 + \sum_{n=1}^{\infty} \left( A_n \cos \frac{n\pi}{L}x + B_n \sin \frac{n\pi}{L}x \right) e^{-\lambda_n \kappa t} \\ u(0, t) = u(L, t) & \end{array}$$

## Question

- ▶ When  $f(x)$  can be written as series? Convergence?
- ▶ If the series of  $f$  converge, will  $u(x, t)$  series converge?
- ▶ If converge, will  $u$  continuous/differentiable/satisfy HE?